

Table 2A

Cancer Incidence in Census Tract 3661 Wayland, MA 1982-1992

Cancer Type	Total				Male				Female			
	Obs	Exp	SIR	95% CI	Obs	Exp	SIR	95% CI	Obs	Exp	SIR	95% CI
Bladder	10	10.1	99	47--482	8	7.1	113	49--223	2	3.0	NC	NC
Breast	39	39.6	98	70--135	1	0.2	NC	NC	38	39.4	97	68--133
Hodgkin's	3	1.9	NC	NC	1	1.0	NC	NC	2	0.9	NC	NC
Kidney	6	5.1	117	43--254	5	3.0	164	53--383	1	2.1	NC	NC
Leukemia	2	4.4	NC	NC	0	2.4	NC	NC	2	2.0	NC	NC
Liver	1	1.1	NC	NC	1	0.7	NC	NC	0	0.3	NC	NC
Lung	37	32.0	116	81--159	17	19.5	87	51--140	20	12.5	160	97--247
NHL	9	7.9	114	52--217	3	3.9	NC	NC	6	4.0	151	55--329
Pancreas	5	5.2	97	31-226	3	2.3	NC	NC	2	2.8	NC	NC
Stomach	3	5.3	NC	NC	2	3.0	NC	NC	1	2.2	NC	NC

Notes:

Expected number of cases presented are rounded to the nearest tenth. SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases Exp = Expected number of cases SIR = Standardized Incidence Ratio NC = Not Calculated

NHL = non-Hodgkin's Lymphoma Lung = Lung and Bronchus 95% CI = 95% Confidence Interval * Indicates statistical significance (P<.05)